#### Superior Urology

146 Court Street South, Thunder Bay, ON P7B2X6 Phone: 807-285-9292 Fax: 855-893-1469

Discharge Instructions Following Cystoscopy or Prostate/Tumor Resection

## <u>Go to an Emergency Department</u> if any of the following occurs:

- You are not able to pass urine
- You are showing signs of infection, a fever of 100.4°F (38°C) or higher, chills, pain with passing urine.
- Frequent urination and/or burning that lasts longer than 2 days.
- Your urine becomes bright or dark red in colour.
- You experience increasing pain or difficulty passing urine

#### **Follow Up Instructions**

Make an appointment to see Dr. \_\_\_\_\_

on/in \_\_\_\_\_.

- Call \_\_\_\_\_\_ to make your appointment.
- If you have any questions, never hesitate to call your doctor.

Other Special Instructions:

Did your doctor tell you that you have developed an infection related to your surgery? If YES, please call the Infection Prevention Control office at LWDH (807) 468-9861 extension 2362 Thank you for helping us ensure guality care.

#### These instructions are intended as a guide.

If you have any questions or concerns, please contact the **LWDH Surgical Daycare Unit** at 807-468-9861 ext 2223,

or the Ministry of Health at 1-888-797-0000, or go to the nearest Emergency Department.

### Lake of the Woods District Hospital

21 Sylvan Street Kenora, Ontario P9N 3W7 Telephone (807) 468-9861

# DISCHARGE INSTRUCTIONS following CYSTOSCOPE



#### The procedure you had was a:

- □ Cystoscopy
- □ Prostate/Tumor Resection

#### Following the Procedure

- You are apt to feel numb after the procedure, the numbness goes away in 1-3 hours. Afterwards, you may feel sore.
- Drinking water helps flush your bladder and reduce irritation. Try to drink 2 glasses of water each hour for the first few hours after the procedure.
- If you've had sedation or general anesthesia, you may be asked to remain in a recovery area until the effects of the medications wear off.
- You may have visible blood in your urine for 24-48 hours. It may also burn when you pass your urine, and/or you may have increased urination for a couple days afterwards.
- If the doctor performed a biopsy, the sample will be sent away for testing.

#### Activity

- Allow your body time to heal. Rest when tired, don't move too quickly, or lift anything heavy until you are feeling better.
- Avoid heavy lifting (over 10 pounds) for \_\_\_\_\_ week(s).
- Try to be active, walking is a great choice for light exercise.
- Many people can return to work within 1 to 3 weeks after surgery. It depends on the type of work you do and how you feel.
- Your doctor will advise you when it's time to return to work.

#### Care at Home

- Take medications as directed if prescribed. (pain control, antibiotics, etc.)
- Drink 8 to 10 glasses of water a day.

- Take a warm bath or place a warm wet washcloth over the opening to the urethra as needed. This will help to ease pain
- Don't delay going to the bathroom, as it can stretch and weaken your bladder. Empty your bladder as soon as you feel the need to.
- Remember it is normal to see some blood in your urine for a few days.

#### What to Watch For (problems that could arise)

- Bleeding
- Infection
- Injury to the bladder and urethra
- Discomfort in the urethra area
- Burning sensation for a short time (common)
- Upset stomach

#### If you have been given anesthesia or sedation

- Do not drive or operate machinery for 24 hours.
- Do not make important decisions or sign any important documents in the next 24 hours. Do not plan any activities that require complex thought or coordination for 24 hours.
- Do not consume alcohol, tranquilizers or sleeping medications for the next 24 hours.
- Sedative and narcotic medicines given during a surgical procedure may linger for hours. You may feel dizzy for several hours after your procedure.
- You must have a responsible adult stay with you tonight.

#### When do I need to call the doctor?

- Large amount of blood in your urine
- Passing large dime-sized clots
- Cloudy urine
- Passing little or no urine
- Abdominal pain and nausea